# **Transport**

## **Question Paper**

Level	Pre U
Subject	Biology
Exam Board	Cambridge International Examinations
Topic	The Life of Plants
Sub Topic	Transport
Booklet	Question Paper

Time Allowed: 26 minutes

Score: /21

Percentage: /100

#### Part - A

Fig. 6.1 is a scanning electronmicrograph of xylem vessels.





Fig. 6.1

(a)	Describe functions.	xylem	vessels,	such	as	those	shown	in	Fig.	6.1,	are	adapted	to their
		 						••••					
		 						••••					
		 						••••					
													[4]

Fig. 6.2 shows the masses of water transpired and the masses of water absorbed by loblolly pine, *Pinus taeda*, and prickly pear cactus, *Opuntia ficus-indica*, over a 24 hour period.



key	
	<ul> <li>loblolly pine transpiration rate</li> <li>loblolly pine water absorption rate</li> </ul>
	<ul> <li>prickly pear transpiration rate</li> <li>prickly pear water absorption rate</li> </ul>

Fig. 6.2

## **Save My Exams! – The Home of Revision**For more awesome GCSE and A level resources, visit us at <a href="https://www.savemyexams.co.uk/">www.savemyexams.co.uk/</a>


(b)	(i)	Describe <b>and</b> explain the patterns of transpiration and water absorption in loblolly pine.
		[4]
	(ii)	Suggest why the patterns of transpiration and water absorption for prickly pear are <b>not</b> the same as for loblolly pine.
		[3]

[Total: 11]

#### Save My Exams! - The Home of Revision

For more awesome GCSE and A level resources, visit us at www.savemyexams.co.uk/

**2** Fig. 23.1 is a photomicrograph of the lower epidermis of the leaf of an oleander, *Nerium oleander*. Fig. 23.2 is a photomicrograph of the lower epidermis of the leaf of a privet, *Ligustrum vulgare*. Both photomicrographs are to the same scale.






Fig. 23.1

Fig. 23.2

(a) State two ways, visible in Fig. 23.1 and Fig. 23.2, in which the epidermis of oleander differs from the epidermis of privet. In each case explain how oleander is adapted to survive severe drought conditions.

1	
2	
	[4]

The concentration of three ions, potassium, chloride and phosphate, were determined in guard cells of closed and open stomata. Fig. 23.3 shows these concentrations measured in arbitrary units, which are the same for all three ions.




Fig. 23.3

(b) Suggest a possible mechanism, that can be supported by the data in Fig. 23.3, to account

for the changes in stomatal aperture.
[4]

### **Save My Exams! – The Home of Revision**For more awesome GCSE and A level resources, visit us at <a href="https://www.savemyexams.co.uk/">www.savemyexams.co.uk/</a>

(c)	Many plant cells have cytoplasmic connections (plasmodesmata) between neighbouring cells, but these are absent from guard cells.
	Explain how this helps guard cells function efficiently.
	[2]
	[Total: 10]