

## Chapter 5 Integration


May/June 2002

6 Let  $f(x) = \frac{4x}{(3x+1)(x+1)^2}$ .

(i) Express  $f(x)$  in partial fractions. [5]

(ii) Hence show that  $\int_0^1 f(x) dx = 1 - \ln 2$ . [5]

10



The function  $f$  is defined by  $f(x) = (\ln x)^2$  for  $x > 0$ . The diagram shows a sketch of the graph of  $y = f(x)$ . The minimum point of the graph is  $A$ . The point  $B$  has  $x$ -coordinate  $e$ .

(i) State the  $x$ -coordinate of  $A$ . [1]

(ii) Show that  $f''(x) = 0$  at  $B$ . [4]

(iii) Use the substitution  $x = e^u$  to show that the area of the region bounded by the  $x$ -axis, the line  $x = e$ , and the part of the curve between  $A$  and  $B$  is given by

$$\int_0^1 u^2 e^u du. \quad [3]$$

(iv) Hence, or otherwise, find the exact value of this area. [3]

Oct/Nov 2002

2 Find the exact value of  $\int_1^2 x \ln x dx$ . [4]

May/June 2003

2 Find the exact value of  $\int_0^1 xe^{2x} dx$ . [4]

10 (i) Prove the identity


$$\cot x - \cot 2x \equiv \operatorname{cosec} 2x. \quad [3]$$

(ii) Show that  $\int_{\frac{1}{6}\pi}^{\frac{1}{4}\pi} \cot x \, dx = \frac{1}{2} \ln 2.$  [3]

(iii) Find the exact value of  $\int_{\frac{1}{6}\pi}^{\frac{1}{4}\pi} \operatorname{cosec} 2x \, dx,$  giving your answer in the form  $a \ln b.$  [4]

Oct/Nov 2003

6



The diagram shows the curve  $y = (3 - x)e^{-2x}$  and its minimum point  $M.$  The curve intersects the  $x$ -axis at  $A$  and the  $y$ -axis at  $B.$

(i) Calculate the  $x$ -coordinate of  $M.$  [4]

(ii) Find the area of the region bounded by  $OA, OB$  and the curve, giving your answer in terms of  $e.$  [5]

8 Let  $f(x) = \frac{x^3 - x - 2}{(x - 1)(x^2 + 1)}.$

(i) Express  $f(x)$  in the form

$$A + \frac{B}{x - 1} + \frac{Cx + D}{x^2 + 1},$$

where  $A, B, C$  and  $D$  are constants. [5]

(ii) Hence show that  $\int_2^3 f(x) \, dx = 1.$  [4]

May/June 2004


- 5 (i) Prove the identity

$$\sin^2 \theta \cos^2 \theta \equiv \frac{1}{8}(1 - \cos 4\theta). \quad [3]$$

- (ii) Hence find the exact value of

$$\int_0^{\frac{1}{3}\pi} \sin^2 \theta \cos^2 \theta \, d\theta. \quad [3]$$

10




The diagram shows the curve  $y = \frac{\ln x}{x^2}$  and its maximum point  $M$ . The curve cuts the  $x$ -axis at  $A$ .

- (i) Write down the  $x$ -coordinate of  $A$ . [1]  
(ii) Find the exact coordinates of  $M$ . [5]  
(iii) Use integration by parts to find the exact area of the shaded region enclosed by the curve, the  $x$ -axis and the line  $x = e$ . [5]

Oct/Nov 2004

7



The diagram shows the curve  $y = x^2 e^{-\frac{1}{2}x}$ .

- (i) Find the  $x$ -coordinate of  $M$ , the maximum point of the curve. [4]  
(ii) Find the area of the shaded region enclosed by the curve, the  $x$ -axis and the line  $x = 1$ , giving your answer in terms of  $e$ . [5]

- 8 An appropriate form for expressing  $\frac{3x}{(x+1)(x-2)}$  in partial fractions is

$$\frac{A}{x+1} + \frac{B}{x-2},$$

where  $A$  and  $B$  are constants.

- (a) Without evaluating any constants, state appropriate forms for expressing the following in partial fractions:

(i)  $\frac{4x}{(x+4)(x^2+3)},$  [1]

(ii)  $\frac{2x+1}{(x-2)(x+2)^2}.$  [2]

(b) Show that  $\int_3^4 \frac{3x}{(x+1)(x-2)} dx = \ln 5.$  [6]

May/June 2005

- 4 (i) Use the substitution  $x = \tan \theta$  to show that

$$\int \frac{1-x^2}{(1+x^2)^2} dx = \int \cos 2\theta d\theta. \quad [4]$$

- (ii) Hence find the value of

$$\int_0^1 \frac{1-x^2}{(1+x^2)^2} dx. \quad [3]$$

- 8 (i) Using partial fractions, find


$$\int \frac{1}{y(4-y)} dy. \quad [4]$$

- (ii) Given that  $y = 1$  when  $x = 0$ , solve the differential equation

$$\frac{dy}{dx} = y(4-y),$$

obtaining an expression for  $y$  in terms of  $x$ . [4]

- (iii) State what happens to the value of  $y$  if  $x$  becomes very large and positive. [1]



The diagram shows part of the curve  $y = \frac{x}{x^2 + 1}$  and its maximum point  $M$ . The shaded region  $R$  is bounded by the curve and by the lines  $y = 0$  and  $x = p$ .

- (i) Calculate the  $x$ -coordinate of  $M$ . [4]
- (ii) Find the area of  $R$  in terms of  $p$ . [3]
- (iii) Hence calculate the value of  $p$  for which the area of  $R$  is 1, giving your answer correct to 3 significant figures. [2]

Oct/Nov 2005


- 6 (i) Use the substitution  $x = \sin^2 \theta$  to show that

$$\int \sqrt{\left(\frac{x}{1-x}\right)} dx = \int 2 \sin^2 \theta d\theta. \quad [4]$$

- (ii) Hence find the exact value of

$$\int_0^{\frac{1}{4}} \sqrt{\left(\frac{x}{1-x}\right)} dx. \quad [4]$$

May/June 2006



The diagram shows a sketch of the curve  $y = x^{\frac{1}{2}} \ln x$  and its minimum point  $M$ . The curve cuts the  $x$ -axis at the point  $(1, 0)$ .


- (i) Find the exact value of the  $x$ -coordinate of  $M$ . [4]
- (ii) Use integration by parts to find the area of the shaded region enclosed by the curve, the  $x$ -axis and the line  $x = 4$ . Give your answer correct to 2 decimal places. [5]

8 Let  $f(x) = \frac{7x+4}{(2x+1)(x+1)^2}$ .

(i) Express  $f(x)$  in partial fractions. [5]

(ii) Hence show that  $\int_0^2 f(x) dx = 2 + \ln \frac{5}{3}$ . [5]

10



The diagram shows the curve  $y = x \cos 2x$  for  $0 \leq x \leq \frac{1}{4}\pi$ . The point  $M$  is a maximum point.

(i) Show that the  $x$ -coordinate of  $M$  satisfies the equation  $1 = 2x \tan 2x$ . [3]

(ii) The equation in part (i) can be rearranged in the form  $x = \frac{1}{2} \tan^{-1}\left(\frac{1}{2x}\right)$ . Use the iterative formula

$$x_{n+1} = \frac{1}{2} \tan^{-1}\left(\frac{1}{2x_n}\right),$$

with initial value  $x_1 = 0.4$ , to calculate the  $x$ -coordinate of  $M$  correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

(iii) Use integration by parts to find the exact area of the region enclosed between the curve and the  $x$ -axis from 0 to  $\frac{1}{4}\pi$ . [5]

May/June 2007

5 (i) Express  $\cos \theta + (\sqrt{3}) \sin \theta$  in the form  $R \cos(\theta - \alpha)$ , where  $R > 0$  and  $0 < \alpha < \frac{1}{2}\pi$ , giving the exact values of  $R$  and  $\alpha$ . [3]

(ii) Hence show that  $\int_0^{\frac{1}{2}\pi} \frac{1}{(\cos \theta + (\sqrt{3}) \sin \theta)^2} d\theta = \frac{1}{\sqrt{3}}$ . [4]

7 Let  $I = \int_1^4 \frac{1}{x(4-\sqrt{x})} dx$ .

(i) Use the substitution  $u = \sqrt{x}$  to show that  $I = \int_1^2 \frac{2}{u(4-u)} du$ . [3]

(ii) Hence show that  $I = \frac{1}{2} \ln 3$ . [6]

Oct/Nov 2007

1 Find the exact value of the constant  $k$  for which  $\int_1^k \frac{1}{2x-1} dx = 1$ . [4]

3 Use integration by parts to show that

$$\int_2^4 \ln x dx = 6 \ln 2 - 2. \quad [4]$$


May/June 2008

7 Let  $f(x) \equiv \frac{x^2 + 3x + 3}{(x+1)(x+3)}$ .

(i) Express  $f(x)$  in partial fractions. [5]

(ii) Hence show that  $\int_0^3 f(x) dx = 3 - \frac{1}{2} \ln 2$ . [4]

9



The diagram shows the curve  $y = e^{-\frac{1}{2}x} \sqrt{1+2x}$  and its maximum point  $M$ . The shaded region between the curve and the axes is denoted by  $R$ .

(i) Find the  $x$ -coordinate of  $M$ . [4]

(ii) Find by integration the volume of the solid obtained when  $R$  is rotated completely about the  $x$ -axis. Give your answer in terms of  $\pi$  and  $e$ . [6]

Oct/Nov 2008

9 The constant  $a$  is such that  $\int_0^a x e^{\frac{1}{2}x} dx = 6$ .

(i) Show that  $a$  satisfies the equation

$$x = 2 + e^{-\frac{1}{2}x}. \quad [5]$$


(ii) By sketching a suitable pair of graphs, show that this equation has only one root. [2]

(iii) Verify by calculation that this root lies between 2 and 2.5. [2]

(iv) Use an iterative formula based on the equation in part (i) to calculate the value of  $a$  correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

May/June 2009

10



The diagram shows the curve  $y = x^2 \sqrt{1 - x^2}$  for  $x \geq 0$  and its maximum point  $M$ .

(i) Find the exact value of the  $x$ -coordinate of  $M$ . [4]

(ii) Show, by means of the substitution  $x = \sin \theta$ , that the area  $A$  of the shaded region between the curve and the  $x$ -axis is given by

$$A = \frac{1}{4} \int_0^{\frac{1}{2}\pi} \sin^2 2\theta d\theta. \quad [3]$$

(iii) Hence obtain the exact value of  $A$ . [4]

Oct/Nov 2009/31


5 (i) Prove the identity  $\cos 4\theta - 4 \cos 2\theta + 3 \equiv 8 \sin^4 \theta$ . [4]

(ii) Using this result find, in simplified form, the exact value of

$$\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \sin^4 \theta d\theta. \quad [4]$$

7





The diagram shows the curve  $y = \frac{\ln x}{\sqrt{x}}$  and its maximum point  $M$ . The curve cuts the  $x$ -axis at the point  $A$ .

- (i) State the coordinates of  $A$ . [1]
- (ii) Find the exact value of the  $x$ -coordinate of  $M$ . [4]
- (iii) Using integration by parts, show that the area of the shaded region bounded by the curve, the  $x$ -axis and the line  $x = 4$  is equal to  $8 \ln 2 - 4$ . [5]

Oct/Nov 2009/32

- 6 (i) Use the substitution  $x = 2 \tan \theta$  to show that

$$\int_0^2 \frac{8}{(4+x^2)^2} dx = \int_0^{\frac{1}{4}\pi} \cos^2 \theta d\theta. \quad [4]$$

- (ii) Hence find the exact value of

$$\int_0^2 \frac{8}{(4+x^2)^2} dx. \quad [4]$$

May/June 2010/31

- 4 (i) Using the expansions of  $\cos(3x - x)$  and  $\cos(3x + x)$ , prove that

$$\frac{1}{2}(\cos 2x - \cos 4x) \equiv \sin 3x \sin x. \quad [3]$$

- (ii) Hence show that

$$\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \sin 3x \sin x dx = \frac{1}{8}\sqrt{3}. \quad [3]$$

8 (i) Express  $\frac{2}{(x+1)(x+3)}$  in partial fractions. [2]

(ii) Using your answer to part (i), show that

$$\left(\frac{2}{(x+1)(x+3)}\right)^2 \equiv \frac{1}{(x+1)^2} - \frac{1}{x+1} + \frac{1}{x+3} + \frac{1}{(x+3)^2}. \quad [2]$$

(iii) Hence show that  $\int_0^1 \frac{4}{(x+1)^2(x+3)^2} dx = \frac{7}{12} - \ln \frac{3}{2}$ . [5]

May/June 2010/32

2 Show that  $\int_0^\pi x^2 \sin x dx = \pi^2 - 4$ . [5]

10 (i) Find the values of the constants  $A$ ,  $B$ ,  $C$  and  $D$  such that


$$\frac{2x^3 - 1}{x^2(2x - 1)} \equiv A + \frac{B}{x} + \frac{C}{x^2} + \frac{D}{2x - 1}. \quad [5]$$

(ii) Hence show that

$$\int_1^2 \frac{2x^3 - 1}{x^2(2x - 1)} dx = \frac{3}{2} + \frac{1}{2} \ln\left(\frac{16}{27}\right). \quad [5]$$

May/June 2010/33

5



The diagram shows the curve  $y = e^{-x} - e^{-2x}$  and its maximum point  $M$ . The  $x$ -coordinate of  $M$  is denoted by  $p$ .

(i) Find the exact value of  $p$ . [4]

(ii) Show that the area of the shaded region bounded by the curve, the  $x$ -axis and the line  $x = p$  is equal to  $\frac{1}{8}$ . [4]

7 (i) Prove the identity  $\cos 3\theta \equiv 4 \cos^3 \theta - 3 \cos \theta$ . [4]

(ii) Using this result, find the exact value of

$$\int_{\frac{1}{3}\pi}^{\frac{1}{2}\pi} \cos^3 \theta d\theta. \quad [4]$$